

Corn production is greatly affected by hybrid selection.

This publication reports the agronomic performance of entries in the 2012 South Dakota performance trials for glyphosate-resistant field corn hybrids.

Major factors in hybrid selection include:

- Yield
- Maturity
- Lodging resistance
- Seed traits

Corn Hybrid Performance Trials Results - Bath

Robert G. Hall | SDSU Extension Agronomist
Kevin K. Kirby | Agricultural Research Manager
Shawn Hawks | Agricultural Research Manager



These performance trials are conducted by the South Dakota Crop Performance Testing program at South Dakota State University. Corn test trial information is listed in table A, performance table footnote references including technology traits and seed treatments are listed in Table B. Mailing addresses for seed company entrants that participated in the 2012 test trials are listed in table C.

General Test Procedures

Seed companies pick one or more of the six test locations where their entries are tested. Entries are placed into "early" or "late" maturity trials. The relative maturity breaks between the early and late tests are 95 days for Bath and South Shore, 100 days for Bancroft and Volga, 105 days for Geddes, and 110 days for Beresford. Hybrids are assigned to trials based on the relative maturity rating reported by the participating seed company. In 2012, results from only five of the test locations are reported. Information from Beresford is not reported because the severe drought at that location prevented the collection of valid data.

Table A. General test information for 2012	
Location:	Glyphosate resistant corn results, Bath, SD
Cooperator:	Gordon and Roger Locken Farms - Bath
Soil Type:	Great Bend silt loam, 0-2% slope
Tillage:	No-till
Fertility-Yield Goal:	200 bushel
Previous Crop:	Soybeans
Row Space:	30 inches
Seeding Population:	30,000/acre
Weed Control:	Pre: Harness Xtra, Post: 1 quart Roundup
Insect Control:	None
Disease Control:	None
Date Seeded:	May 1, 2012

Experimental Procedures

Entries were seeded in three replications, with each hybrid randomly located within each trial. Plots consisted of four 30-inch rows that were 20-feet long, with the center two rows harvested for yield. A Monosem precision row crop planter was used for seeding plots at all locations. In 2012, the precision planter was calibrated to deliver 29,621 seeds per acre, regardless of seed quality and germination percentage. No seeding rate adjustment was made for low germination. Therefore, percent stand is an indication of initial seed quality and the ability of the seed to cope with the production environment from seeding to harvest. Soil type, land preparation and previous crop history, pest control, and fertility yield goal at each test site is outlined in table A. Seedbed preparation was good at all locations. A starter fertilizer of 100 pounds/acre of 37-18-00 was applied 2" below and 2" to the side (2 x 2) of the seed row.

Measures of Performance

Yields are obtained from the South Dakota Crop Performance

Testing Program. Two-year and current yield averages are included where hybrids have been tested for the most recent two-year period.

Yield. Yield values are an average of three replications and are expressed as bushels per acre, adjusted to 15.5% moisture on a dry-matter basis and a bushel weight of 56 pounds. Statistical tests were conducted to determine whether differences obtained were caused by variations in environment or were true hybrid differences. In 2012, the coefficient of variation (CV) values (a measure of experimental error) for yield was quite variable, ranging from a low of 4% in both trials at Bath to a high of 12% and 15% for the early and late trials, respectively, at Geddes. Experimental error may be the result of several factors, including test methods, or factors such as moisture, temperature, soil variations, or agronomic factors like seeding date, reseeding, or seed quality factors—all of which may or may not be controllable in a given year. Clearly, this year, drought

and variable seasonal moisture distribution and/or subsoil moisture conditions, along with above-average temperatures, combined to produce highly variable yields across our South Dakota corn growing regions. The top yield averages occurred at Bath and Bancroft, whereas, the plot grain yields were so low at Beresford the harvest equipment could not determine valid harvest yields.

Grain moisture content.

Moisture content is expressed as the percentage of moisture in the shelled corn at harvest. Hybrids that are above average in yield and can be stored without additional drying are desirable. At harvest, moisture values were determined by the combine moisture meter, which in turn was periodically verified with a Dickey-John GAC II meter.

Use of tables. Check for the "least significant difference" (LSD) value at the bottom of each column of data averages. The LSD value indicates how much a variable such as yield must differ between two hybrids

before there is a significant yield difference. LSD values are given at the bottom of every column where there is significant difference among the averages within the column. If differences among the averages within a column are not significant, the LSD value is reported as “nonsignificant” (NS). The top performance group (TPG) for the performance factors yield (bu/a), bushel weight in (lbs.), harvest

grain moisture (%), lodging below the ear (%), and final stand (%) are indicated in the performance tables by the shaded data values. Evaluate the performance of each hybrid (row) by moving across the table and observing if any of the reported performance values are shaded. The more shaded performance values in each row, the better the hybrid did as a top-performing hybrid for a given test trial.

Performance Trial Results

Note: Evaluate the performance of each hybrid (row) by moving across the table and observing if any of the reported performance values are shaded. The more shaded performance values there are in each row the better the hybrid did as a top-performing hybrid for a given test trial.

No.	Explanation of references
[1]	Entries are listed by Brand/Variety– Entries are sorted by 2-yr then by 2012 yield average.
[2]	Technology traits: Agr. = Agrisure; SmartStax or GenSSX= DowAgrosciences (DAS) or Monsanto Genuity (Gen), respectively; HX1, HXX = Herculex by DAS & Pioneer Hi-Bred, Opt. = Optimum AcreMax1 or Optimum AcreMaxXtra by Pioneer Hi-Bred, YG = Monsanto YieldGard.
[3]	Seed treatments: P/V500 = Poncho/Votivo 500, P/V1250 = Poncho/Votivo1250; CM250 = Cruiser Maxx250, CM1250 = Cruiser Max1250; MHL = Maxim XL, MQ = Maxim Quattro; PPST250 = Pioneer Premium Seed Treatment.
[4]	Brand Relative Maturity (Rel. Mat.)– the relative maturity rating reported by the test trial entrant.
[5]	Lodging Percentage– stalks broken below the ear as a percentage of the final stand; look for low values.
[6]	Final Stand Percentage – the number of standing stalks at harvest as a percentage of the seeding population.
[7]	Least Significant Difference (LSD 0.05) – the difference any two values within a column must be equal to or exceed for the values to be significantly different (0.05 level of probability). If the difference is less than the LSD value the difference between them is nonsignificant (NS).
[8]	Min. TPG-avg.– the minimum column value for yield, bushel weight, and final stand percentage that a hybrid must equal or exceed to be in the top performance group (TPG).
[9]	Max. TPG-avg.– the maximum column value for grain moisture at harvest, lodging percentage, or lodging score that a hybrid must equal or be less than to be in the TPG.
[10]	Coefficient of variation (C.V.)– the percent of experimental error associated with a test trial. Ideally, the CV value for yield is less than 15%. Values less than 5% are less common ; while values of 6-15% are more common. If a value exceeds 15%; the trial had too much error to be valid, so the results are not reported.

Table 1. Bath early maturity Roundup Ready corn hybrid test results, 2011-12, Gordon and Roger Locken Farm.

Brand/Hybrid [1] + Tech. Trait [2] + Seed Trt. [3]	Rel. Mat. [4]	Yield Averages *,#		Other 2012 Averages##			
		2-Yr bu/a	2012 bu/a	Bu. Wt. lb	Grain Moisture %	Ldg. % [5]	Final Stand % [6]
SEEDS/ 2000 2903GTCELL + AgrGT/CB/LL + Acceleron	90	223	218	56	15	1	87
RENK/ RK585VT3P + GenVT3P + Acceleron	95	222	215	58	14	0	94
SEEDS 2000/ 9504VT3P + GenVT3P + Acceleron	95	217	212	59	14	1	89
MASTERS/ CHOICE MCT-493 + Agr3000GT + CM250+MXL	93	212	192	57	13	1	80
G2/ GEN. 5X-795 + HXXRR2 + MQ	95	207	215	58	13	0	91
SEEDS 2000/ 9202 VT2P + OptAMX + MQ	92	207	210	57	13	0	89
WENSMAN/ W 7140VT3PRO + GenVT3P + Acceleron	93	207	206	60	14	1	93
SEEDS 2000 /9503 VT2P + GenVT2P + Acceleron	95	206	204	60	15	1	88
NUTECH/ 5N-9001 + LL + CM250	90	206	188	58	13	0	80
PETERSON/ FARMS 76R92 + GenVT2P + Acceleron	92	204	202	58	13	0	89
WENSMAN/ W 8120VT2RIB + GenVT2P + Acceleron	92	203	206	57	13	0	86
RENK/ RK530VT3P + GenVT3P + Acceleron	94	202	204	60	13	0	90
PIONEER/ 38H08 + HX1RR2 + PPST250	92	201	199	57	12	2	89
CHANNEL/ 192-09VT3P + GenVT3P + P/V500	92	.	220	58	13	0	92
RENK/ RK492VT3P + GenVT3P + Acceleron	93	.	215	60	13	0	91
PETERSON FARMS/ PFS76Z95 + GenVT2P + Acceleron	95	.	213	59	16	1	90
RENK/ RK568VT3P + GenVT3P + Acceleron	95	.	212	59	15	0	93
G2 GEN./ 5X-193 + HXXRR2 + P/V1250	93	.	211	58	13	1	88
DEKALB/ DKC43-48 + GenVT3P + P/V500	93	.	210	58	13	1	94
EPLEY/ E9505RR + RR2 + MQ	95	.	209	60	14	0	89
PROSEED/ 1292VT2P + LL + Acceleron	92	.	205	58	13	0	89
DEKALB/ DKC43-10 + GenVT2P + P/V500	93	.	204	58	13	0	87
WENSMAN/ W 8184VT3PRO + GenVT3P + Acceleron	95	.	201	58	15	2	88
RENK/ RK434VT3P + GenVT3P + Acceleron	92	.	200	59	13	0	85
PROSEED/ 1295SS + LL + Acceleron	95	.	197	60	13	0	93
G2 GEN./ 5X-9402 + HXXRR2 + P/V1250	94	.	197	59	13	0	83
PROSEED/ D99031116 + LL + CM250+MQ	90	.	194	58	14	0	75
PETERSON FARMS/ PFS75T93 + GenVT2P + Acceleron	93	.	194	60	13	1	88
PIONEER/ P8906HR + HX1RR2 + PPST250	89	.	194	59	13	1	92
Trial avg.:	93	209	205	58	14	1	88
High avg.:	95	223	220	60	16	2	94
Low avg.:	89	201	188	56	12	0	75
[7] LSD(.05):		NS**	14	1	1	1	5
[8] Min.TPG value:		201	206	59	.	.	89
[9] Max.TPG value:		.	.	.	13	1	.
[10] Coef. of var.:		4	4	1	4	187	3
No. entries:	29	13	29	29	29	29	29

NOTE: Table references [1-10] are explained in table B.

* Entries are listed by Brand/Hybrid and sorted by 2-yr then by 2012 yield average.

Adjusted to 15.5% moisture.

** NS indicates differences between column values are nonsignificant.

Shaded column values are included in the top-performance group (TPG).

Table 2. Bath late maturity Roundup Ready corn hybrid test results, 2011-12, Gordon and Roger Locken Farm.

Brand/Hybrid [1] + Tech. Trait [2] + Seed Trt. [3]	Rel. Mat. [4]	Yield Averages *,#		Other 2012 Averages##			
		2-Yr bu/a	2012 bu/a	Bu. Wt. lb	Grain Moisture %	Ldg. % [3]	Final Stand % [6]
NUTECH/ 5N-001 + Agr3000GT + MQ	100	224	217	56	16	0	81
PIONEER/ 36V53 + HX1RR2 + PPST250	102	222	219	56	15	0	91
DEKALB/ DKC50-66 + YGVTWR2 + P/V500	100	221	219	58	13	0	91
DAIRYLAND/ ST-9399 + Agr3000GT + CM250	99	221	216	57	15	0	89
CHANNEL/ 196-06VT3P + GenVT3P + P/V500	96	221	212	59	15	0	89
SEEDS 2000/ 9902 VTP3111 + Agr311 + Acceleron	98	213	206	57	14	1	93
DEKALB/ DKC48-12 + GenSSX + P/V500	98	208	206	57	13	0	82
WENSMAN/ W 7268VT3 + YGVTWR2 + Acceleron	96	205	208	57	16	0	92
RENK/ RK580VT3 + GenSSX + Acceleron	98	203	184	58	15	0	78
PIONEER/ P0193HR + HX1RR2 + PPST250	101	.	228	54	16	2	89
PIONEER/ P0062XR + HXXRR2 + PPST250	100	.	227	54	15	8	96
G2 GEN./ 5X-0004 + HXXRR2 + P/V1250	100	.	225	54	15	1	90
DEKALB/ DKC49-30 + GenVT3P + P/V500	99	.	222	59	14	0	95
PIONEER/ P9834HR + HX1RR2 + PPST250	98	.	220	55	14	0	81
G2 GEN./ 5Z-198 + OptAMX + P/V1250	98	.	220	55	12	0	86
DAIRYLAND/ DS9501SSX + DAS Smart Stax + Activa Compl.	101	.	219	56	13	0	92
WENSMAN/ W 7290VT3PRO + GenVT3P + Acceleron	99	.	218	58	14	0	96
WENSMAN/ W 9288VT3PRO + GenVT3P + Acceleron	98	.	218	58	15	0	91
SEEDS 2000/ 3011 G3 + Agr3000GT + Acceleron	101	.	218	55	15	0	81
PIONEER/ P9917AM1 + OptAM1 + PPST250	99	.	214	58	14	0	88
WENSMAN/ W 7320VT3PRO + GenVT3P + Acceleron	101	.	208	58	15	1	90
G2 GEN./ 5X-895 + HXXRR2 + MQ	97	.	207	57	13	0	88
NUTECH/ 5N-0103 + Agr3000GT + CM250	101	.	205	56	15	0	87
G2 GEN./ 5H-502 + HX1RR2 + MQ	102	.	204	56	16	0	79
PIONEER/ P9675AMX + OptAMX + PPST250	96	.	203	58	13	0	85
RENK/ RK578VT3P + GenVT3P + Acceleron	96	.	195	58	13	0	89
DEKALB/ DKC46-20 + GenVT3P + P/V500	96	.	194	60	15	0	87
DAIRYLAND/ DS9303SSX + DAS Smart Stax + Activa Compl.	103	.	194	55	15	1	88
RENK/ RK576VT3P + GenVT3P + Acceleron	97	.	191	58	13	0	89
NUTECH/ 5B-798 + AgrGT/CB/LL + CM250	98	.	181	58	16	1	78
DAIRYLAND/ DS9402SSX + DAS Smart Stax + Activa Compl.	102	.	172	54	14	1	82
Trial avg.:	99	215	209	57	14	1	88
High avg.:	103	224	228	60	16	8	96
Low avg.:	96	203	172	54	12	0	78
[5] LSD(.05):		NS**	12	2	1	1	6
[6] Min.TPG value:		203	216	58	.	.	90
[7] Max.TPG value:		.	.	.	13	1	.
[8] Coef. of var.:		3	4	2	5	143	4
No. entries:	31	9	31	31	31	31	31

NOTE: Table references [1-10] are explained in table B.

* Entries are listed by Brand/Hybrid and sorted by 2-yr then by 2012 yield average.

Adjusted to 15.5% moisture.

** NS indicates differences between column values are nonsignificant.

Shaded column values are included in the top-performance group (TPG).

Early maturity trial, Table 1.

The test-trial yield averages were **209 bu/a** for two years and **205 bu/a** in 2012. Hybrids that yielded **201 bu/a** or more for two years and **206 bu/a** or more for 2012 qualified for the top-performance-group (TPG) for yield. There was no significant difference in yield average among the hybrids tested for the two year period. Hybrids had to differ in yield average by **14 bu/a** in 2012 to be significantly different. In 2012, bushel weights averaged **58 lbs.**, grain moisture averaged **14%**, lodging percentage averaged **1%**, and final stand percentage averaged **88%**. In order for a hybrid to be in the TPG for these factors, it had to average **59 lbs.** or more in bushel weight, **13%** or less in grain moisture, **1%** or less in lodging

percentage, and **89%** or more for final stand percentage. In order for a hybrid variable to differ from one hybrid compared to another, their difference had to equal **14 bu/a** or more for yield in 2012, **1 lb.** or more in bushel weight, **1%** or less in grain moisture, **1%** or less in lodging percentage and **5%** or more in the final stand percentage.

Late maturity trial, Table 2.

The test-trial yield averages were **215 bu/a** for two years and **209 bu/a** in 2012. Hybrids with yield averages of **203 bu/a** or more for two years and **216 bu/a** or more for 2012 qualified for the TPG. There was no significant difference in yield average among the hybrids tested for the two year period. Hybrids had to differ

in yield average by **12 bu/a** in 2012 to be significantly different. In 2012, bushel weights averaged **57 lbs.**, grain moisture averaged **14%**, lodging percentage averaged **1%**, and final stand percentage averaged **88%**. In order for a hybrid to be in the TPG for these factors, it had to average **58 lbs.** or more in bushel weight, **13%** or less in grain moisture, **1%** or less in lodging percentage, and **90%** or more for final stand percentage. In order for a hybrid variable to differ from one hybrid compared to another, their difference had to equal **12 bu/a** or more for yield in 2012, **2 lb.** or more in bushel weight, **1%** or more in grain moisture, **1%** or more in lodging percentage and **6%** or more in the final stand percentage.

Table C. Mailing addresses for seed entries in the 2012 corn hybrid trials and listed by seed brand name	
Seed brand	Seed company mailing address
Channel	Channelbio Corp., Box 277, Laurel, NE 68745
Dairyland	PO Box 958, West Bend, WI 53095
Dekalb	46040 SD Hwy 44, Chancellor, SD 57015
Epley Bros.	Epley Bros. Hybrids Inc., 22494 Yale Ave., Shell Rock, IA 50670
G-2 Genetics	G-2 Genetics, 2321 North Loop Drive, Suite 230, Ames, IA 50010
Hoegemeyer	Hoegemeyer Hybrids, 1755 Hoegemeyer Road, Hooper, NE 68031
Masters Choice	3010 St. Rt. 146 E, Anna, IL 62906
NuTech	Nutech Seed, LLC, 2321 North Loop Drive, Suite 230, Ames, IA 50010
Petersen Farms	3104 164th Ave. SE, Harwood, ND 58042
Pioneer	Pioneer Hi-Bred International, 151 Saint Andrews Court-Suite 910, Mankato, MN 56001
ProSeed	705 E. Brewster, Harvey, ND 58341
Renk	6809 Wilburn Road, Sun Prairie, WI 53590
Seeds 2000	Seeds 2000, 115 N 3rd St., Breckenridge, MN 56520
Wensman	Wensman Seed Co., PO Box 190, Wadena, MN 56482



South Dakota State University, South Dakota counties, and U.S. Department of Agriculture cooperating. South Dakota State University is an Affirmative Action/Equal Opportunity Employer and offers all benefits, services, education, and employment opportunities without regard for race, color, creed, religion, national origin, ancestry, citizenship, age, gender, sexual orientation, disability, or Vietnam Era veteran status.