

## BEST MANAGEMENT PRACTICES

### CHAPTER 26



## Starter, Banding, and Broadcasting Phosphorus Fertilizer for Profitable Corn Production

Anthony Bly (Anthony.Bly@sdstate.edu), Graig Reicks (Graig.Reicks@sdstate.edu), and Ron Gelderman (retired professor, SDSU Plant Science Dept.)

The phosphorus (P) application methods used in corn production fields can impact yield and net profitability. The primary benefit from banding P is that it concentrates the P in a small zone near the plant, which can improve P availability. The objectives of this chapter are to: 1) discuss different P application methods, 2) summarize research comparing P application methods, 3) review the effect of band distance from the seed, 4) discuss safe amounts of P-containing fertilizers that can be placed with the seed, and 5) consider strengths and weaknesses of the application methods.

### Introduction

Plant-available P moves slowly through the soil by diffusion to the root. Diffusion is the movement of ions from an area with high concentration to an area with low concentration. Factors that influence P diffusion are soil texture, water content, bulk density, pH, temperature and the distance between higher soil P concentrations zones such as bands and the root. P diffusion rates are higher for finer textured soil (clay), higher soil water content, lower soil bulk density, soil pH between 5.0 and 6.0, higher soil temperatures, and shorter distances from root to soil zones with higher P concentration (Bhadoria et al., 1991; Grierson et al., 1999; Olsen and Watanabe, 1963; Schachtman et al., 1998). Crops with finer root systems or those that have been colonized by arbuscular mycorrhizal fungi (AMF) may have higher P uptake efficiencies (Bittman et al, 2006). Adopting no-till practices that encourage fungi growth and development can encourage AMF colonization.




Figure 26.1 P applied as a broadcast or banded application. (Courtesy of Colorado State University, Clay et al., 2009)

Phosphorus fertilizer application methods include broadcast, banded, or band-applied with the seed (Fig. 26.1). Liquid or dry fertilizers are used with any of these placement options. Starter fertilizers, placed at planting, can be located to the side of the row or with the seed (pop-up). Usually, only a portion of the total recommended P rate for optimum corn yield is placed as a starter, unless the recommended P rate is low enough to enable full rate application. The balance of the P recommendation that is not applied as a starter needs to be applied with another method.

Concentrating fertilizer P in a band often improves P availability as there is less opportunity for the fertilizer P to be tied up in the soil, especially at very low or high soil pH. Rates can sometimes be reduced by one-third or more for band-applied P. However, reducing rates can result in a decline of soil test P over time, which can reduce yield potential (Chapters 23 and 29).

Phosphorus can be applied in a variety of forms ranging from liquid to solid products (Chapter 28) and the application technique will depend on your goals, your soil test value, and tillage and equipment options. Tillage system and equipment availability play an important role in determining what choice a corn grower makes when considering P placement. Equipping a very large planter with fertilizer application equipment can be less than desirable because the added weight can lead to higher soil compaction. Frequently asked questions regarding P placement for corn:

1. What P application method returns the greatest economic return?
2. What distance from the corn row should P bands be placed?
3. How much fertilizer can be placed with the seed at planting?

### P Application Method Comparisons (band vs. broadcast)

Choosing the most appropriate P application method is complicated by the many factors that influence plant P uptake efficiency. Placing the P in a band near the seed may increase plant P uptake efficiency because it concentrates P in a zone easily reached by early root growth. In addition, P diffusion from the band to the root is high due to a large concentration gradient when compared with broadcast P application. However, when the bulk soil test P is high, the benefits of banded P are reduced. Therefore, knowing soil test P level is important when determining placement options. The optimal balance between P banding and broadcast application is difficult to achieve (Barber, 1974, 1985).

Soil fertility researchers conducted research studies at several land-grant universities between 1970 and 2015 that compared banded and broadcast P applications (Fig. 26.2).

This research suggested that the benefit from banding P increased with yield potential. For example, at yield less than 100 bu/acre banding and broadcasting P had similar yields, whereas at 200 bu/acre, banding P had a 7.4 bu/acre yield advantage over broadcasting P. When the P rate was considered, the yield advantage from banding was higher when the P application rate was  $\leq 40$  lbs  $P_2O_5$ /acre (Fig. 26.3). Based on these findings, banding is recommended when yields are greater than 150 bu/acre, P rates  $< 40$  lbs  $P_2O_5$ /acre, and soil test P is




Figure 26.2 Band and broadcast P corn grain yield comparison from P placement literature review. (1970-2010)




Figure 26.3 Band and broadcast P relative grain yield comparisons across P rates taken from literature review. (1970-2010)

at or below the medium category.

These recommendations are contrary to findings from Iowa, where banded and broadcast treatments had similar yields (Mallarino et al., 1999; 2004). Differences between South Dakota and Iowa may be attributed to differences in rainfall, length of growing season, and tillage.

Could higher soil moisture conditions in South Dakota no-till cornfields result in similar conclusions between banded and broadcast P? Phosphorus placement research projects were conducted in South Dakota between 1998 and 1999 at nine sites located in corn grower no-till fields. Phosphorus fertilizer was either placed as starter (2x2 band) or broadcast at 40 lbs  $P_2O_5$ /acre. Statistical comparison between the two treatments showed that site year was significant, which is indicative of a wide range of yield environments used in the project. The banded P treatment (134.1 bu/acre) grain yield average was greater than broadcast (130.8 bu/acre), but not statistically significant. A relative grain yield comparison of broadcast and banded P across Olsen P soil test levels showed that banded P resulted in greater yield at the 4-6 ppm soil test range and similar yields when the soil test value was  $\geq 10$  ppm (Fig. 26.4).




Figure 26.4 Relative no-till corn grain yield of broadcast and banded P comparisons at 9 locations in SD from 1998 to 1999.




Figure 26.5 P band distance from corn seed furrow influence on relative grain yield at 4 locations in eastern South Dakota during 2004 and 2005.

### Distance of P Band from the Seed

Phosphorus band distance from the crop root is another factor influencing P uptake efficiency. Few research studies have investigated the optimal P band distance because it requires specialized research equipment. Research conducted in 2004 and 2005 near Beresford, SD, and Brookings, SD, evaluated the effect of P band distance from the seed furrow on grain yield. This research showed that the greatest yield increase occurred when the P band was located 2 inches from the seed (Fig. 26.5).

### Seed-placed Phosphorus

In some situation, placing the P fertilizer with the seed can improve P uptake efficiency. Triple super phosphate (0-46-0) was a very common P fertilizer material in the past and was very safe for seed application. However, for various reasons, this material is not produced to a great extent by the plant food industry. More recently developed P sources containing nitrogen include ammonium polyphosphate (10-34-0), monoammonium phosphate (MAP, 11-52-0), and diammonium phosphate (DAP, 18-46-0). A number of liquid P sources containing N, P, and K have been developed such as 7-21-7, 4-10-10, 3-18-18 and 9-18-9. In addition, numerous mixed fertilizers have been developed containing secondary and micronutrients as well. A pop-up fertilizer calculator has been developed for determining safe pop-up

| Soil Texture | Soil Moisture at Planting                     |            |           |
|--------------|---|------------|-----------|
|              | Dry   | Borderline | Moist     |
|              | ----- gallons/acre (lbs $P_2O_5$ /acre) ----- |            |           |
| Fine/Medium  | 5.8 (23)                                      | 7.8 (31)   | 11.7 (46) |
| Coarse       | 4.7 (19)                                      | 5.8 (23)   | 7.8 (31)  |

fertilizer rates (Gelderman, 2007). This calculator considers the fertilizer source, soil texture, soil moisture, row width, and risk adversity. The calculator was developed using laboratory method results that were well-correlated to field study data.

This calculator is available from the International Plant Nutrition Institute, [www.ipni.net](http://www.ipni.net) (search for Seed-placed Fertilizer Decision Aid). Table 26.1 was developed from this tool and shows the allowable rates of ammonium polyphosphate fertilizer (APP) with differing soil moisture and textures that can be applied with the corn seed planted in 30-inch rows using a 5% maximum seed germination reduction risk.

### Summary

There are many different techniques that can be used to apply P to soils. Each application approach has strengths and weaknesses (Table 26.2). South Dakota has a frigid, semi-arid environment that may influence plant responses to P compared with other corn growing regions. In our soils and climatic conditions, management practices that encourage early growth may result in higher corn yields. Different outcomes may be found in more temperate environments. This paper contains several key summaries including:

1. Understanding soil phosphorus, the meaning of soil P test levels, and the factors influencing P movement (diffusion) to plant roots are very important for optimal P placement.
2. At lower  $P_2O_5$  application rates (< 40 lbs/acre), band-placement grain yields were higher than broadcast and generally occurred at research sites with soil test P below the medium level.
3. In South Dakota no-till corn research, corn grain yield from fertilizer band (2x2) applications were higher than broadcast applications at soil test (Olsen) P levels less than 10 ppm.
4. P bands placed 4 inches or less from the seed furrow returned the highest grain yield.
5. P placement with the seed is a good management practice. However, special attention needs to be taken to understand how different fertilizer materials may reduce seed germination. The seed-placed fertilizer calculator, [www.ipni.net](http://www.ipni.net), is a good tool to determine pop-up fertilizer limits.

**Table 26.2 Strengths and weaknesses of P application methods.**

|           | Strengths   | Weaknesses  |
|-----------|---|---|
| Pop-up    | <ul style="list-style-type: none"> <li>• Promotes early growth.</li> <li>• Can increase yields if soil test values are low.</li> <li>• Increases P uptake efficiency when soil test P is lower.</li> </ul>  | <ul style="list-style-type: none"> <li>• Can reduce germination if the rate is too high.</li> <li>• Ammonia contained in the fertilizer can reduce germination.</li> <li>• Extra equipment for planter to carry.</li> <li>• Some liquids are more corrosive and can damage equipment over time.</li> <li>• Additional P fertilizer may have to be applied to get recommended P rate.</li> </ul> |
| Broadcast | <ul style="list-style-type: none"> <li>• Can be easily applied.</li> <li>• Will increase the bulk soil P level.</li> <li>• Will increase corn yields.</li> <li>• Could be as effective when compared with banding when rates &gt; 40 lbs <math>P_2O_5</math>/acre.</li> </ul>   | <ul style="list-style-type: none"> <li>• May be less effective than banding for low P rates.</li> <li>• May be less effective when soil test P levels are low.</li> </ul>   |
| Banded    | <ul style="list-style-type: none"> <li>• In high-yield environment can increase yields.</li> <li>• Most effective if the soil P level is in the low to medium level.</li> <li>• Bands should be placed less than 5 inches from the seed.</li> <li>• Increases P uptake efficiency when soil test P is lower.</li> </ul> | <ul style="list-style-type: none"> <li>• May not be effective for low-yield potentials.</li> <li>• May not be practical for high P rates.</li> <li>• Extra equipment for planter to carry.</li> <li>• Some liquids are more corrosive and can damage equipment over time.</li> <li>• Additional P fertilizer may have to be applied to get the recommended rate.</li> </ul>                     |

## References and Additional Information

- Barber, S.A. 1976. Efficient fertilizer use. P. 13-29. In F.L. Peterson (ed.) *Agronomic research for food*. Am. Soc. of Agron., Madison WS.
- Barber, S.A., and J.L. Kovar. 1985. Review – Principles of Applying Phosphorus Fertilizer for Greatest Efficiency. *J. of Fertilizer Issues*, 2:3: 91-94.
- Bhadoria, P.B.S., J. Kaselowsky, N. Claassen, and A. Jungk. 1991. Phosphate Diffusion Coefficients in Soil as Affected by Bulk Density and Water Content. *J Plnt Nut Soil Sci*, 154:53-57.
- Bittman, S., C.G. Kowalenko, D.E. Hunt, T.A. Forge, and X. Wu. 2006. Starter Phosphorus and Broadcast Nutrients on Corn with Contrasting Colonization by Mycorrhizae. *Agron.J.* 98:394-401.
- Bordoli J.M., and A.P. Mallarino. 1998. Deep and shallow banding of phosphorus and potassium alternatives to broadcast fertilization for no-till corn. *Agron. J.* 90:27-33.
- Buah S., T.A. Polito, and R. Killorn. 2000. No-Tillage Corn Response to Placement of Fertilizer Nitrogen, Phosphorus, and Potassium. *Commun. Soil Sci. Plant Anal.* 32: 3121-3133.
- Cihacek L.J., D.L. Mulvaney, R.A. Olson, L.F. Welch and R.A. Wiese. 1974. Phosphate placement for corn in chisel and moldboard plow systems. *Agron. J.* 66:665-668.
- Clay, D.E., S.A. Clay, and K. Reistma. 2009. *South Dakota Corn Best Management Practices*. 2009. South Dakota State Univeristy, Brookings, SD. .
- Eckert D.J. and J.W. Johnson. 1985. Phosphorus Fertilization in No-tillage Corn Production. *Agron. J.* 77:789-792.
- Fixen P.E., B.G. Farber, and M. Vivehanandan. 1987. Phosphorus management as influenced by tillage systems in eastern South Dakota 1984-1986. *Ag. Exp. Station and Plant Science Dept. SDSU. Annual Progress Report. Plant Science Pamphlet TB99.*
- Gelderman, R.H. 2007. Fertilizer placement with seed: A decision aid. In: *Proceedings of the North Central Extension-Industry Soil Fertility Conference*, Des Moines. IA. Nov. 14-15, 2007. Vol 23. Int. Plant Nutrition Inst. Brookings, SD. P. 46-54.
- Gelderman R., A. Bly, and A. Huer. 2010. Influence of phosphorus placement and rate on no-till corn, NE farm 2010. 2010 Soil and Water Research. *Ag. Exp. Station and Plant Science Dept. SDSU. Annual Progress Report. Plant Science Pamphlet No. 45. Soil PR 10-18.*
- Gelderman R., A. Bly, and A. Huer. 2009. Influence of phosphorus placement and rate on no-till corn, NE farm 2009. 2009 Soil and Water Research. *Ag. Exp. Station and Plant Science Dept. SDSU. Annual Progress Report. Plant Science Pamphlet No. 42. Soil PR 09-11.*
- Gelderman R., J. Gerwing, B. Berg, and A. Bly. 1999. Phosphorus rate and placement effects on tilled corn and soybean rotation. 1999 Soil and Water Research. *Ag. Exp. Station and Plant Science Dept. SDSU. Annual Progress Report. Plant Science Pamphlet TB 99. Soil PR 99-8.*
- Gelderman R., J. Gerwing, and R. Berg. 2001. Long-term phosphorus management for no-till corn and soybean in the northern cornbelt. 2001 Soil and Water Research. *Ag. Exp. Station and Plant Science Dept. SDSU. Annual Progress Report. Plant Science Pamphlet No. 9. Soil PR 00-6.*
- Gelderman, R.H., and L. Ulvestad. 2010. A Summary of soil test results. (July 2009-June 2010). 2010 Soil and Water Research. *Ag. Exp. Station and Plant Science Dept. SDSU. Annual Progress Report. Plant Science Pamphlet No. 45. Soil PR 10-5.*



- Gelderman, R., J. Gerwing, and A. Bly. 1999. Determination of critical soil test K level and starter P needs for no-till corn in the northern corn-belt. 1999 Soil and Water Research. Ag. Exp. Station and Plant Science Dept. SDSU. Annual Progress Report. Plant Science Pamphlet TB 99. Soil PR 99-7.
- Gerwing J., and R. Gelderman. 2005. Fertilizer Recommendations Guide. South Dakota State University, Cooperative Extension Service and U.S. Department of Agriculture. Plant Science, EC-750.
- Grierson P.F., N.B. Comerford, and E.J. Jokela. 1999. Phosphorus mineralization and microbial biomass in a Florida Spodosol: effects of water potential, temperature and fertilizer application. *Biol Fertil Soils* 28:244-252.
- Howard D.D., M.E. Essington, and J. Logan. 2002. Long-Term broadcast and banded phosphorus fertilization of corn produced using two tillage systems. *Agron. J.* 94:51-56.
- Mallarino A.P., J.M. Bordoli, and R. Borges. 1999. Phosphorus and potassium placement effects on early growth and nutrient uptake of no-till corn and relationship with grain yield. *Agron. J.* 91:37-45.
- Mallarino A.P., and D. Haden. 2004. Phosphorus and potassium fertilizer placement for corn and soybeans managed with no-till and chisel-disk tillage. Iowa State University Farm Progress Reports. Paper 1367.
- Olsen S.R., and F. S. Watanabe. 1963. Diffusion of phosphorus as related to soil texture and plant uptake. *Soil Sci. Soc. Am. J.* 1963. 27:648-653.
- Rehm G.W. 1986. Effect of phosphorus placement on early growth, yield and phosphorus absorption by irrigated corn. *Journal of Fertilizer Issues.* 3:1 pp. 12-17. Rehm G.W. 1996. It's not just starter and broadcast anymore. University of Minnesota. Department of Soil, Water, and Climate. pp. 74-83.
- Reitsma, K., D.E. Clay, C.G. Carlson, A. Bly, and G. Reicks. 2016. Estimating yield goals and nitrogen, phosphorus, and potassium recommendations. Chapter 23 In Clay, D.E., C.G. Carlson, S.A. Clay, and E. Byamukama (eds). *iGrow Corn: Best Management Practices*. South Dakota State University.
- Riedell W.E., D.L. Beck, and T.E. Schumacher. 2000. Corn response to fertilizer placement treatments in an irrigated no-till system. *Agron. J.* 92:316-320.
- Schachtman D.P., R.J. Reid, and S.M. Ayling. 1998. Phosphorus uptake by plants: from soil to cell. *Plant Physiol.* 116: 447-453.

## Acknowledgements

Support for this document was provided by South Dakota State University, SDSU Extension, South Dakota Corn Utilization Council and International Plant Nutrition Institute (IPNI).



Bly, A, G. Reicks, and R. Gelderman. 2015. Chapter 26: Starter, Banding, and Broadcasting Phosphorus Fertilizer for Profitable Corn Production. In Clay, D.E., C.G. Carlson, S.A. Clay, and E. Byamukama (eds). iGrow Corn: Best Management Practices. South Dakota State University.

*The preceding is presented for informational purposes only. SDSU does not endorse the services, methods or products described herein, and makes no representations or warranties of any kind regarding them.*

*In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident.*

*Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotope, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English.*

*To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, found online at [http://www.ascr.usda.gov/complaint\\_filing\\_cust.html](http://www.ascr.usda.gov/complaint_filing_cust.html) and at any USDA office or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by:*

(1) mail: U.S. Department of Agriculture  
Office of the Assistant Secretary for Civil Rights  
1400 Independence Avenue, SW  
Washington, D.C. 20250-9410;

(2) fax: (202) 690-7442; or

(3) email: [program.intake@usda.gov](mailto:program.intake@usda.gov).

*USDA is an equal opportunity provider, employer, and lender.*